

# A New Folk Theorem in OLG Games

Chihiro Morooka (諸岡千裕)

March 11, 2020

## Abstract

In this paper, we analyze the model of general  $n$ -person overlapping generations games. It is assumed that players strictly discount the future at a common rate and that the dimension of feasible one-shot payoffs with individual rationality is equal to  $n$ . In contrast to previous researches, this study shows that players can obtain payoffs outside the feasible set of one-shot games.

*JEL Classification Codes:* C72; C73.

*Keywords:* Overlapping Generations Games; Intertemporal Trade; Discount Factor; Folk Theorem; Cubic Hull.

## 1 Introduction

Overlapping generations (OLG) games have been widely studied to investigate economic organizations whose members change over time. In OLG games, players in the same generation interact for a sufficiently long time, and then are replaced sequentially by successors in the next generation. Kandori (1992) first proved the folk theorem in general  $n$ -person OLG games, which stated that the pair of players in each generation can obtain any payoffs in  $V^*$ , the convex combination of feasible one-shot payoffs with individual rationality. Following him, Smith (1992) proved several versions of folk theorems with stronger results.

In the analyses of Kandori (1992), Smith (1992), and subsequent studies on OLG games, the characterization of equilibrium set of payoffs is restrictive. They showed that players can obtain any payoffs in  $V^*$  without discounting, and that the result is robust against low discounting. However, this does not mean that players cannot attain payoffs outside  $V^*$ : when they discount the future, intertemporal trades of payoffs using the difference of time preferences can be possible.

Based on this issue, we recently have shown in Morooka (2020) that in 2-person OLG games with discounting, players can obtain payoffs

outside  $V^*$ . In the present paper, we extend the result of Morooka (2020) to general  $n$ -person OLG games. Our result shows that under the full dimensionality of  $V^*$ , any payoffs in the cubic hull of  $V^*$ , that is, any payoffs in the smallest  $n$ -dimensional cube that contains  $V^*$ , are obtainable by the pair of players in each generation.

In order to get the intuition of our result, we consider the one-shot game illustrated in Figure 1. In our model of overlapping generations, each player lives for  $3T$  periods, and his first and second opponents are replaced by corresponding new players in  $T$  and  $2T$  periods, respectively, after his entrance, as depicted in Figure 2. Players are assumed to discount the future at a common rate. When each player takes  $Y$  in the first  $T$  periods, and  $X$  in the next  $2T$  periods, an equilibrium is formed with an average payoff of  $\frac{3(1-\delta^T)}{1-\delta^{3T}}$  which is sufficiently close to 3 as the player's continuation payoffs diminish as periods pass. As a result, players' payoffs in each generation are almost  $(3, 3, 3)$ , which Pareto-dominate  $(1, 1, 1)$  and are outside  $V^*$ .

The main difference between Morooka (2020) and this work is in rewarding players after punishments. In 2-person games, mutual min-maxing is available for punishment against deviations by one player. When there are 3 or more players in one-shot games, they may want to quit punishment because their payoffs during punishment are less than their minimax payoffs. Players must be rewarded with appropriate payoffs after the punishment as an incentive, which requires the full dimensionality of feasible one-shot payoffs with individual rationality.

In order to obtain this result, players' strict discount is essential. In our study, the rate of discounting has two different roles, which can be seen as a trade-off. First, players must be so patient that do not deviate, the argument of which is seen everywhere in the literature on repeated games. At this point, we do not want players to discount the future. The second role, which is our original one, is that it must diminish the payoffs in players' later days. If players do not discount the future, they fail the intertemporal trades of payoffs.

It must also be noted that there is an order in the choice of parameters; we must fix the discount first, and then choose players' lifespan, depending on the discount. This is because, as the discount rate approaches 1, we need a longer lifespan in order for players' continuation payoffs after they become older to diminish sufficiently. This order is inverse to that of the non-uniform OLG folk theorem in Smith (1992), where he first fixes players' lifespan, allowing players to punish devi-

		$A_2$	
		$A_3: X$	
		$X$	$Y$
$A_1$	$X$	1, 1, 1	<u>0, 3, 0</u>
	$Y$	<u>3, 0, 0</u>	<u>0</u>

		$A_2$	
		$A_3: Y$	
		$X$	$Y$
$A_1$	$X$	<u>0, 0, 3</u>	<u>0</u>
	$Y$	<u>0</u>	<u>0</u>

Figure 1: Stage game 1

	Gen.1			Gen.2			Gen.3			...		
$A_1$	1~3T			3T+1~6T			6T+1~9T			...		
$A_2$	(1~T)	T+1~4T			4T+1~7T			7T+1~10T			...	
$A_3$	(1~2T)			2T+1~5T			5T+1~8T			8T+1~11T		...

Figure 2: Structure of overlapping generations in the case of  $n=3$

ations strictly under no discount, and then chooses the discount with which the punishments are still available.

In relation to infinitely repeated games, our logic is compared to that of Lehrer and Pauzner (1999). They studied two-person infinitely repeated games with different discounting between players, and showed that players can obtain equilibrium payoffs outside the set of feasible one-shot payoffs  $V$ . In their model, the player who evaluates future payoffs at a higher rate gives payoffs to his opponent with less patience first, and is rewarded later. This allows players to obtain higher payoffs outside  $V$ . In Lehrer and Pauzner (1999), however, there is an asymmetry of payoffs between players. In our model, on the other hand, all players' payoffs can be raised at no cost.

We define the model and show the results in Section 2 and Section 3 concludes.

## 2 Model and Results

Let  $N=\{1,2,\dots,n\}$  for  $n \geq 2^1$  represent the set of players in a one-shot game. For  $i \in N$ , let  $A_i$  be the set of  $i$ 's actions and let  $A=\prod_{i \in N} A_i$ . Let  $g_i$  be the one-shot payoff function of  $i$ . Let  $G=G(N, A, g)$  be the one-shot game considered. Without loss of generality, we assume that each player's minimax payoff in  $G$  is 0. Let  $V=co(g(A))$  be the set of feasible payoffs in  $G$ . The set of feasible and individually rational payoffs is then defined as  $V^*=V \cap R^{n+}$ . We assume the full dimensionality, that is,  $\dim V^*=n$ . We also assume that the origin is included in  $V^*$ .

<sup>1</sup>Although our result holds in the case of  $n=2$ , it is included in the stronger result of Morooka (2020) without full dimensionality condition.

In order to rigorously characterize the set of equilibrium payoffs in our result, consider the following value  $r_i$  for  $i \in N$ :

$$r_i = \max_{v \in V^*} v_i.$$

This value is the best feasible and individually rational payoff for player  $i$ . We then define the *open cubic hull* of  $V^*$ , say  $C(V^*)$ , as follows:

$$C(V^*) = \prod_{i \in N} (0, r_i).$$

For example, in the game shown in Figure 1,  $r_1=r_2=r_3=3$  and  $C(V^*)=(0, 3)^3$ .

We now turn to construct our model of OLG games,  $OLG(G; \delta, T)$ , as follows. Let  $T$  be a natural number. Each player lives for  $nT$  periods: the player of  $k$ 'th generation with action  $A_i$  lives between period  $(k-1)nT + (i-1)T + 1$  and  $knT + (i-1)T$ . Exclusively, 0'th generation with action  $A_i$  for  $i \neq 1$  lives between period 1 and  $(i-1)T$ . The case of  $n=3$  is shown in Figure 2.

In our model, each player's future payoffs are discounted at a common rate  $\delta \leq 1$ <sup>2</sup>. It is assumed that players can use the public randomizing device each period and that the monitoring is perfect. Let  $g_i(t)$  be the one-shot payoff of a player with the action set  $A_i$  in period  $t$ . When the sequence of actions  $\{a_t\}_{t=1}^{nT}$  is played throughout a player's life with action set  $A_i$  (except in the case of players in 0'th generation), his average payoff is as follows:<sup>3</sup>:

$$\frac{1}{\sum_{t=1}^{nT} \delta^{t-1}} \sum_{t=1}^{nT} \delta^{t-1} g_i(a_t).$$

The following result holds.

**Theorem 1 (The Folk Theorem in discounted  $n$ -person OLG Games).** For every  $v \in C(V^*)$ , and for every  $\epsilon > 0$ , there is  $\delta_0 \in (0, 1)$  such that if  $\delta \in [\delta_0, 1)$ , then for every  $T$  sufficiently large, there is a subgame perfect equilibrium in  $OLG(G; \delta, T)$ , where the pair of players in each generation<sup>4</sup> obtains average payoffs  $\epsilon$ -close to  $v$ . This vector in particular need not be feasible.

<sup>2</sup>Although our result does not hold with  $\delta=1$ , it is convenient to consider this case in order to construct the strict punishments under no discount which is still available with  $\delta < 1$ .

<sup>3</sup>For players in 0'th generation, please replace  $nT$  by their lifespans

<sup>4</sup>Players in 0'th generation with shorter lifespans are excluded from this result.

Before the formal proof, we see in detail that in the OLG model of the following stage game in Figure 3, each player can obtain  $v_i=5$  within the error of  $\epsilon=0.1$  in an equilibrium, with  $\delta=0.95$ . In the Pareto frontier of this game, the sum of payoffs is 13, which is less than  $\sum_i v_i=15$ . In this game,  $Y$  is the strongly dominant action for all players. However, in order to generate efficient (resp. minimaxing) outcome, two players must take  $X$  (resp.  $M$ ) enduring payoff  $-1$ , which is less than the minimax value. Moreover, the only one Nash profile  $(Y, Y, Y)$  is not only Pareto-inferior but also gives considerably high payoffs to players, so it cannot be used both on the path and for punishments. It is assumed that in each period, the nature draws one integer from 1 to 16 with equal probability, and that players commonly observe it. Consider the following strategy.

If there is no deviation, players play the following path strategy:

- When a player exits, he takes  $Y$  and his opponents take  $X$ .

- In all the other periods, players play the action which gives 5 to the youngest player and gives 4 to the other players. This payoff is generated when players play  $(Y, X, X)$  (which yields 15 to the youngest player and  $-1$  to his opponents) with probability  $3/8$ ,  $(X, Y, X)$  with probability  $5/16$ , and  $(X, X, Y)$  with probability  $5/16$ .

If someone deviates, players play one from the following punishments. The punishments are also applied to the deviations from themselves.

- (Punishment 1) If there are at least three periods left between the next period and the oldest player's exit, or if a player deviates at his opponent's exit, the deviator plays  $Y$  and his opponents play  $M$  in the next period. Then, in the following period, players play the action which gives 1 to the deviator and gives 3 to his opponents. The payoffs  $(1, 3, 3)$  are generated when players play  $(Y, X, X)$  with probability  $1/8$ ,  $(X, Y, X)$  with probability  $1/4$ ,  $(X, X, Y)$  with probability  $1/4$ , and  $(M, M, M)$  with probability  $3/8$ . After that, they return to the path.

- (Punishment 2) If the oldest player deviates and there are 2 periods

left before his exit, he plays  $Y$  and his opponents play  $M$  until he exits. After that, players return to the path.

·(Punishment 3) If the oldest player deviates 1 period before his exit, he plays  $Y$  and his opponents play  $M$  in the next period. After that, players return to the path.

·(Punishment 4) If a younger player deviates and there are at most 2 periods left before the oldest player's exit, players play  $(Y, Y, Y)$  until the exit of oldest. After the exit, players play the action which gives 3 to them for 12 periods. Then, they return to the path. The payoffs  $(3, 3, 3)$  are generated when players play  $(Y, X, X)$ ,  $(X, Y, X)$ ,  $(X, X, Y)$ , and  $(M, M, M)$  with equal probability.

We show that the above strategy forms an equilibrium. First, observe that players' minimum one-shot payoff is  $-1$  and that their maximum one-shot gain from deviations is 1, by the construction of equilibrium strategy. Consider the deviation of one player when there are at least three periods left between the next period and the oldest player's exit, or when a player deviates at his opponent's exit. If he does not deviate, his payoff in the current period and next two periods is at least  $-1 + 3\delta(1 + \delta)$ . If he deviates, it reduces to at most  $1 + \delta^2$  by Punishment 1. Because  $\delta=0.95$ , the former is greater than the latter.

Second, consider the oldest player's deviation right before his exit. When there are 3 periods between the current period and his exit, his payoff from the current period on is at least  $-1 + 3\delta + 15\delta^2$  without deviations. If he deviates, it reduces to at most 1 by Punishment 2. When there are 2 periods before his exit, his payoff without deviations is at least  $-1 + 15\delta$ , whereas it reduces to at most 1 by Punishment 3 if he deviates.

Third, consider a younger player's deviation right before the oldest player's exit. When there are 3 periods between the current period and the oldest player's exit, his payoff from the current period and the 12'th period after the exit is at least  $-1 + 3\delta - \delta^2 + 4\delta^3(\sum_{t=1}^{12} \delta^{t-1})$  if he does not deviate. If he deviates, it reduces to at most  $1 + 4.2\delta(1 + \delta) + 3\delta^3(\sum_{t=1}^{12} \delta^{t-1})$  by Punishment 4. When there are 2 periods between the current period and the exit, the payoff is at least  $-1 - \delta + 4\delta^2(\sum_{t=1}^{12} \delta^{t-1})$  if he does not deviate. If he deviates, it reduces to at most  $1 + 4.2\delta + 3\delta^2(\sum_{t=1}^{12} \delta^{t-1})$ .

Finally, take  $T \geq 51$ . Then, players' payoff is almost 4.9.

Now, we proceed to formally prove the result. We generalize the punishments against the deviations described above.

Proof. Fix  $v=(v_1^1, \dots, v_n^n) \in C(V^*)$  and  $\epsilon > 0$ . By the definition of  $C(V^*)$ , there exists an  $(n-1)$ -dimensional vector  $v_{-i}^i$  for  $i \in N$  which satisfies  $v^i=(v_i^i, v_{-i}^i) \in \text{int}.V^*$ . Let  $a^i$  be a (correlated) profile with one-shot payoffs  $v^i$ . Because  $\dim V^*=n$  and of the assumption that the origin is in  $V^*$ , there exists a (correlated) profile  $c$  and  $d^i$  with one-shot payoffs  $g(c) \in V^*$ ,  $g(d^i) \in V^*$  satisfying  $g_i(d^i) < g_i(d^j)$  (no dev. from minimax  $j$ ),  $g_i(d^i) < g_i(c)$  (no dev. from  $c$ ) and  $g_i(c) < v_i^j$  (no dev. to  $c$ ) for  $j \neq i$ <sup>5</sup>. Players take  $c$  instead of  $a^j$  to punish the player with  $A_i$  after he becomes the older player, when he deviated right before the exit of his opponent with  $A_j$ , whereas  $d^i$  is played as a "reward" after the punishment against him when he deviated far before the currently oldest player's exit. We denote the minimaxing profile against  $i$ , the profile which gives the maximum payoff to  $i$ , and a one-shot Nash profile, as  $m^i$ ,  $b^i$ , and  $e$ , respectively.

The game starts with playing  $a^1$  in period 1. The parameters  $Q$ ,  $M$ ,  $S$ , and  $P$  are natural numbers and determined later in this order for which the following strategy profile is subgame perfect. The profile consists of six steps, and we need the observability of mixed actions to work all the steps well, except Step 2. We also need PRD for Steps 1 and 4 in order to let players play  $a^i$  and  $c$ .

When there is no deviation, players play the following path strategy. Suppose that the youngest (resp. oldest) player has the action set  $A_i$  (resp.  $A_j$ ) for  $i \in N$  and  $j \neq i$ .

Step 1. When there are at least  $S$  periods left between the next period and the oldest player's exit, players play  $a^i$ .

Step 2. When there are at most  $S$  periods left between the current period and the oldest player's exit, players play  $b^j$ .

---

<sup>5</sup>We can indeed find such  $c$  and  $(d^i)_{i \in N}$ . To see this, choose any  $i \in N$ . There exists a vector on the line connecting the origin and  $v^i$ , each of whose components is less than  $\min_{(i,j)} v_i^j$ . Because of the assumption that the origin is in  $V^*$ , we can choose  $c$  as a pair of actions giving such a vector. Then, because of full-dimensionality, we can choose a vector in the interior of  $V^*$  such that only the  $i$ 'th component is less than  $g_i(c)$ , with remaining other components being the same as  $g(c)$ . We can choose  $d^i$  as giving such a vector.

After someone unilaterally deviates, players play the following punishments. The punishments are also applied to the deviations from themselves. Right after the punishments, players return to the path. Suppose that the deviator has the action set  $A_i$  for  $i \in N$ .

Step 3. When there are at least  $Q+M+S$  periods left between the next period and the oldest player's exit, or when a player deviates at the period of his opponent's exit, players play  $m^i$  for  $Q$  periods, and then play  $d^i$  for  $M$  periods.

Step 4. When the deviation is by a younger player and there are at most  $Q+M+S$  periods left between the current period and the oldest opponent's exit, players play  $e$  until the opponent's exit. After that, players play  $c$  for  $P$  periods.

Step 5. When the deviation is by the oldest player and there are at most  $Q+M+S$  periods left between the current period and his exit, players play  $m^i$  until he exits.

When players play this strategy profile,  $a^i$  is played for  $T-S$  periods on the equilibrium path when the player with  $A_i$  is the youngest, which guarantees that his average payoff is almost equal to  $v_i^i$  under the positive rate of discounting and sufficiently long lifespan. In the remainder of this section, it is proved that the above strategy profile forms a subgame perfect equilibrium for  $\delta$  sufficiently close to 1 when we choose the length of punishments appropriately. The methodology is to provide strictly positive penalties for any unilateral deviation from each step when  $\delta=1$ . Then, by continuity of payoffs, there is some  $\delta_0 \in (0, 1)$  for which all penalties are still positive for  $\delta \in [\delta_0, 1)$ . For each step, the "worst-case scenario" is concerned, where the incentive to deviate is greatest.

We define the following variables:  $u_i = \min_{j \neq i} \{v_i^i, v_i^j, g_i(c), g_i(d^j)\}$  is the worst one-shot payoff for players in Steps 1 and 4.  $\beta$  is the maximum one-shot payoff for players.  $\underline{\beta}$  is the least payoff among players' maximum one-shot payoffs, and  $\omega$  is the minimum one-shot payoff for players. Suppose that the deviator has the action set  $A_i$ .



• Deviations from Steps 1, 3, and 4 when there are at least  $Q+M+S$  periods left between the next period and the oldest player's exit:  
 First, suppose that no player is to be minimaxed in the current period. The deviator gets at most  $\beta$  immediately, and at most 0 in the following  $Q$  periods by Step 3. When he does not deviate, his payoff through these  $1+Q$  periods is at least  $g_i(d^i)$ . We can avoid the deviation when we choose  $Q$  satisfying the following equations for all  $i \in N$ :

$$(1 + Q)g_i(d^i) > \beta.$$

Second, suppose that his opponent is to be minimaxed in the current period. When he deviates in this period, he gets at most  $\beta$  immediately, and at most 0 in the following  $Q$  periods, and then at most  $g_i(d^i)$  in the next  $M$  periods. When he does not deviate, his payoff through these  $1+Q+M$  periods is at least  $Q\omega+(M+1)u_i$ . We can avoid the deviation when we choose  $M$  satisfying the following equations for  $i \in N$ :

$$Q\omega + (M + 1)u_i > \beta + Mg_i(d^i).$$

• Deviations by the oldest player when there are at most  $Q+M+S$  periods left between the current period and his exit:  
 Suppose that the oldest player deviates from the beginning of Step 3 and there are  $Q+M+S - 1$  periods left between the next period and his exit. He gets at most  $\beta$  immediately, and at most 0 in the following  $Q+M+S-1$  periods because of Step 5. When he does not deviate, his payoff through these  $Q+M+S$  periods is at least  $(Q+M)\omega+S\underline{\beta}$ . We can avoid the deviation when we choose  $S$  satisfying the following equations for  $i \in N$ :

$$(Q + M)\omega + S\underline{\beta} > \beta.$$

• Deviations by a younger player when there are at most  $Q+M+S$  periods left between the current period and the oldest player's exit:  
 Suppose that the player deviates from the beginning of Step 3 and there are  $Q+M+S - 1$  periods left between the next period and the exit of oldest player with  $A_j$ . He gets at most  $\beta$  immediately and  $(Q+M+S - 1)g_i(e)$  in the following  $Q+M+S - 1$  periods. After the

oldest player's exit, his payoff in the following  $P$  periods is reduced to  $Pg_i(c)$  because of Step 4. When he does not deviate, he gets at least  $(Q+M+S)\omega$  in the following  $Q+M+S$  periods and at least  $Pv_i^j$  after the entry of younger opponent. We can avoid the deviation when we choose  $P$  satisfying the following equations for  $i \in N$  and  $j \neq i$ :

$$(Q + M + S)\omega + Pv_i^j > \beta + (Q + M + S - 1)g_i(e) + Pg_i(c).$$

Therefore, deviations from any steps strictly decrease deviators' payoffs under no discount, which means that the above strategy profile forms a subgame perfect equilibrium for some  $\delta_0 \in (0, 1)$  and  $\delta \in [\delta_0, 1)$ . Fix such a  $\delta$ . Note that the parameters  $Q$ ,  $M$ ,  $S$ , and  $P$  are independent of the players' lifespan. We then choose  $T_0$  which satisfies the following equations:

$$T_0 > P + Q + M + S,$$

$$\frac{1}{1 - \delta^{nT_0}} \{(1 - \delta^{T_0-S})v_i^i + (\delta^{T_0-S} - \delta^{nT_0})\omega\} > v_i^i - \epsilon, \text{ and}$$

$$\frac{1}{1 - \delta^{nT_0}} \{(1 - \delta^{T_0-S})v_i^i + (\delta^{T_0-S} - \delta^{nT_0})\beta\} < v_i^i + \epsilon \text{ for } i \in N.$$

In the last two equations, each LHS is the minimum and maximum of each player's average payoff, respectively. These values converge to  $v_i^i$  for any  $\delta < 1$  when  $T_0 \rightarrow \infty$ . Set  $T \geq T_0$ . Then, the average payoff of each player with  $A_i$  is almost equal to  $v_i^i$  within the distance of  $\epsilon$ .  $\square$

### 3 Conclusion

In this paper, we have analyzed the model of  $n$ -person OLG games. Under the condition of full dimensionality, we have seen that players can obtain average payoffs outside the convex hull of one-shot payoffs. One remaining question is how the result will change when the stage game does not have the full dimensionality. Even in such a case, players may still be able to obtain payoffs outside the feasible set of one-shot payoffs. Consider the pure coordination game in Figure 4. Clearly, the dimension of this game is 1. For example, when players play  $Y$  in the first  $T$  periods and  $X$  in the following  $2T$  periods, one player attains almost 1 and his opponents attain almost 0. As a result,

<i>A3: X</i>	<i>X</i>	<i>Y</i>	<i>M</i>
<i>X</i>	-1, -1, -1	-1, 15, -1	-1, -1, -1
<i>Y</i>	15, -1, -1	1, 1, -1	1, -1, -1
<i>M</i>	-1, -1, -1	-1, 1, -1	-1, -1, -1

<i>A3: Y</i>	<i>X</i>	<i>Y</i>	<i>M</i>
<i>X</i>	-1, -1, 15	-1, 1, 1	-1, -1, 1
<i>Y</i>	1, -1, 1	4.2, 4.2, 4.2	1, -1, 1
<i>M</i>	-1, -1, 1	-1, 1, 1	-1, -1, 0

<i>A3: M</i>	<i>X</i>	<i>Y</i>	<i>M</i>
<i>X</i>	-1, -1, -1	-1, 1, -1	-1, -1, -1
<i>Y</i>	1, -1, -1	1, 1, -1	0, -1, -1
<i>M</i>	-1, -1, -1	-1, 0, -1	-1, -1, -1

Figure 3: Stage game 2

<i>A3: X</i>	<i>X</i>	<i>Y</i>
<i>X</i>	0	
<i>Y</i>		

<i>A3: Y</i>	<i>X</i>	<i>Y</i>
<i>X</i>	0	0
<i>Y</i>	0	1

Figure 4: Pure coordination game

payoffs in the cube  $(0, 1)^3$  are attainable in equilibria of the OLG game. We will study further whether players can still attain payoffs outside one-shot feasible payoffs in other games without full dimensionality.

## Acknowledgments

I would like to thank Michihiro Kandori, Akihiko Matsui, Tadashi Sekiguchi and Yosuke Yasuda for their helpful comments and suggestions. I would also like to acknowledge various useful ideas from audiences at my presentation during the Japanese Economic Association 2019 autumn meeting at Kobe University.

## References

- Kandori, M. (1992) “Repeated Games Played by Overlapping Generations of Players”, *Rev. of Econ. Stud.* 59, 81-92.
- Lehrer, E., and Pauzner, A. (1999) “Repeated Games with Differential Time Preferences”, *Econometrica* 67, 393-412.
- Morooka, C. (2020) “Equilibrium Payoffs in Overlapping Generations Games with Discounting”, Revised and Resubmitted to *Int. Jour. of Game Theory*.
- Smith, L. (1992) “Folk Theorems in Overlapping Generations Games”, *Games and Econ. Behav.* 4, 426-449.